

M

EDCO FORUM[®]

PRESENTING INNOVATIVE PRODUCTS & SERVICES TO HEALTHCARE PROFESSIONALS

VOLUME 11 NUMBER 22

MAY 2004

REPRINT

TRILYTE[™] WITH FLAVOR PACKS: A LOWER-SODIUM, HIGH-PEG LAVAGE WITH FIVE FLAVOR PACKS

Schwarz Pharma (Milwaukee, WI), recently introduced new TriLyte[™] with flavor packs, a lower-sodium, high-PEG lavage. TriLyte with flavor packs offers five flavors packaged with just one bottle, which may improve patient compliance and promote a clear colonoscopy. It is the only lower-sodium PEG lavage with five flavor packs for true patient choice. TriLyte with flavor packs has proven to be safe and effective

in a wide range of patients. As with all polyethylene glycol (PEG) lavages, nausea, abdominal fullness, and bloating are the most common adverse reactions, occurring in up to 50% of patients. Oral medication administered within 1 hour of the start of administration of TriLyte with flavor packs may be flushed from the gastrointestinal tract and not absorbed.

Introducing the One, the Only Tastebuddies!

The Five Great Tastes of TriLyte[™] with Flavor Packs — a Reduced-Salt PEG Lavage

Available in:
Lemon Lime
Orange
Pineapple
Cherry
Citrus Berry

- Five Great Flavors—Just One Bottle
- Patient Flavor Preference May Enhance Compliance
- Featuring a Lower-Sodium PEG Lavage Formulation
- Proven Safe and Effective in a Wide Range of Patients

TriLyte[™] with flavor packs

(PEG-3350, sodium chloride,
sodium bicarbonate and
potassium chloride for oral solution)

As with all polyethylene glycol (PEG) lavages, nausea, abdominal fullness and bloating are the most common adverse reactions, occurring in up to 50% of patients. Oral medication administered within 1 hour of the start of administration of TriLyte[™] with flavor packs may be flushed from the gastrointestinal tract and not absorbed.

Brief Summary: Before prescribing, please see the complete Prescribing Information.

Indications and Usage: TriLyte[™] with flavor packs is indicated for bowel cleansing prior to colonoscopy. **Contraindications:** TriLyte[™] with flavor packs is contraindicated in patients known to be hypersensitive to any of the components. TriLyte[™] with flavor packs is contraindicated in patients with ileus, gastrointestinal obstruction, gastric retention, bowel perforation, toxic colitis or toxic megacolon. **Warnings:** The flavor packs are for use only with the accompanying 4 liter bottles. No additional ingredients, e.g. flavorings, should be added to the solution. TriLyte[™] with flavor packs should be used with caution in patients with severe ulcerative colitis. Use of TriLyte[™] with flavor packs in children younger than 2 years of age should be carefully monitored for occurrence of possible hyponatremia, as the solution has no caloric substrate. Dehydration has been reported in 1 child and hypokalemia has been reported in 3 children. **Precautions: General:** Patients with ingested egg shells, unconscious or semiconscious patients, and patients prone to regurgitation or aspiration should be observed during the administration of TriLyte[™] with flavor packs, especially if it is administered via nasogastric tube. If a patient experiences severe bloating, distention or abdominal pain, administration should be slowed or temporarily discontinued until the symptoms abate. If gastrointestinal obstruction or perforation is suspected, appropriate studies should be performed to rule out these conditions before administration of TriLyte[™] with flavor packs. **Adverse Reactions:** Nausea, abdominal fullness and bloating are the most common adverse reactions (occurring in up to 50% of patients) to administration of TriLyte[™] with flavor packs. Abdominal cramps, vomiting and anal irritation occur less frequently. These adverse reactions are transient and subside rapidly. Isolated cases of urticaria, rhinitis, dermatitis and (rarely) anaphylactic reaction have been reported which may represent allergic reactions. **Oral Administration: Adults:** At a rate of 240 mL (8 oz.) every 10 minutes, until the rectal effluent is clear or 4 liters are consumed. **Pediatric Patients: (aged 6 months or greater):** At a rate of 25 mL/kg/hour, until the rectal effluent is clear or 4 liters are consumed. Rapid drinking of each portion is preferred to drinking small amounts continuously.

© 2004 Schwarz Pharma, Inc., Milwaukee, WI 53201 USA

Z11246 05/04

TriLyte[™] with flavor packs

(PEG-3350, sodium chloride,
sodium bicarbonate and
potassium chloride for oral solution)

For more information concerning TriLyte, call Schwarz Pharma at 1-800-558-5114; contact a Schwarz Pharma representative at DDW, booth #2835; visit the product Web site at www.trilyte.com, or the company's Web site at www.schwarzpharma.com.

Medco Forum[®] is a registered trademark of Medco Communications LLC Copyright © 2004 Medco Communications LLC. On editorial matters or to request additional copies, telephone (303) 674-9607. Any reproduction, in whole or in part, without express written permission of publisher is prohibited. The information and statements directed to the products discussed herein are based solely on information and statements received from manufacturers and/or distributors thereof. The publishers do not warrant and assume no liability for the accuracy of the information contained herein. The manufacturers and/or distributors should be contacted for any and all information that the reader may desire. Send inquiries or comments regarding this publication to: Medco Communications LLC, 87 Oak Way, Evergreen, CO 80439.